

**COLLECTIVE ORDER CONVERGENCE AND
COLLECTIVELY QUALIFIED SETS OF OPERATORS**E.YU. EMELYANOV *Communicated by E.M. RUDOY*

Abstract: Collective versions of order convergences and corresponding types of collectively qualified sets of operators in vector lattices are investigated. It is proved that collectively order to norm bounded sets are bounded in the operator norm and collectively order continuous sets are collectively order bounded.

Keywords: vector lattice, order convergence, collectively qualified set of operators.

1 Introduction

The study of collectively compact sets of operators between normed spaces was initiated by Anselone and Palmer [6] (for recent advances see [9, 10, 12]). In many cases it is necessary to deal with “uniform” or “collective” properties of a set of operators between vector lattices (shortly, VLs) regarding the order convergence in their domains. Whereas the corresponding technique is rather simple for the norm convergence, the case of order convergence in VLs requires an additional attention. The present note is

EMELYANOV, E.YU., COLLECTIVE ORDER CONVERGENCE AND COLLECTIVELY QUALIFIED SETS OF OPERATORS.

© 2025 EMELYANOV E.YU.

The research was carried out within the framework of the state contract of the Sobolev Institute of Mathematics (project no. RWNF-2026-0022).

Received September, 5, 2024, Published April, 14, 2026.

devoted to investigation of collective order convergence and its applications to collectively qualified sets of operators in VLs.

We abbreviate a normed space (normed lattice, Banach lattice) by NS (NL, BL). Throughout the text, vector spaces are real, operators are linear, symbol $\mathcal{L}(X, Y)$ stands for the space of operators between vector spaces X to Y , symbol $x_\alpha \downarrow 0$ for a decreasing net in a VL such that $\inf_\alpha x_\alpha = 0$, symbol $\text{sol}(A)$ for the solid hull $\bigcup_{a \in A} [-|a|, |a|]$ of a subset A of a VL E , and $\mathcal{L}(E, F)$ ($\mathcal{L}_+(E, F)$, $\mathcal{L}_{ob}(E, F)$, $\mathcal{L}_{oc}(E, F)$) for the space of linear (resp., positive, order bounded, order continuous) operators between VLs E and F .

We shall use the following modes of convergence in vector lattices.

Definition 1. A net (x_α) in a vector lattice E

- a) *order converges* to $x \in E$ (briefly, $x_\alpha \overset{o}{\rightarrow} x$) if there exists a net (g_β) in E such that $g_\beta \downarrow 0$ and, for each β , there is an α_β with $|x_\alpha - x| \leq g_\beta$ for all $\alpha \geq \alpha_\beta$.
- b) *unbounded order converges* to $x \in E$ (briefly, $x_\alpha \overset{uo}{\rightarrow} x$) if $|x_\alpha - x| \wedge u \overset{o}{\rightarrow} 0$ for every $u \in E_+$.
- c) *relative uniform converges* to $x \in E$ (briefly, $x_\alpha \overset{ru}{\rightarrow} x$) if, for some $u \in E_+$ there exists an increasing sequence (α_n) of indices such that $|x_\alpha - x| \leq \frac{1}{n}u$ for all $\alpha \geq \alpha_n$.

An operator T from a VL E to a VL F is called *o-(uo-, ru-)-continuous* if $Tx_\alpha \overset{o}{\rightarrow} 0$ ($Tx_\alpha \overset{uo}{\rightarrow} 0$, $Tx_\alpha \overset{ru}{\rightarrow} 0$) whenever $x_\alpha \overset{o}{\rightarrow} 0$ (resp., $x_\alpha \overset{uo}{\rightarrow} 0$, $x_\alpha \overset{ru}{\rightarrow} 0$).

The following notions of “collective convergence” are useful in working with families of nets indexed by the same directed set.

Definition 2. Let $\mathcal{B} = \{(x_\alpha^b)_{\alpha \in A}\}_{b \in B}$ be a set of nets indexed by a directed set A in a vector lattice E . We say that \mathcal{B}

- a) *collective o-converges* to 0 (briefly, $\mathcal{B} \overset{c-o}{\rightarrow} 0$) if there exists a net $g_\beta \downarrow 0$ such that, for each β , there is an α_β with $|x_\alpha^b| \leq g_\beta$ for $\alpha \geq \alpha_\beta$ and $b \in B$.
- b) *collective uo-converges* to 0 (briefly, $\mathcal{B} \overset{c-uo}{\rightarrow} 0$) if $\{(|x_\alpha^b| \wedge u)_{\alpha \in A}\}_{b \in B} \overset{c-o}{\rightarrow} 0$ for every $u \in E_+$.
- c) *collective ru-converges* to 0 (briefly, $\mathcal{B} \overset{c-ru}{\rightarrow} 0$) if, for some $u \in E_+$, there exists an increasing sequence (α_n) of indices such that $|x_\alpha^b| \leq \frac{1}{n}u$ for $\alpha \geq \alpha_n$ and $b \in B$.

For the basic theory of VLs we refer to [3, 4, 13, 14].

2 Main results

2.1. Preliminaries.. We begin with the following auxiliary properties of c-o, c-uo, and c-ru convergences.

Proposition 1. Let $\mathcal{B} = \{(x_\alpha^b)\}_{b \in B}$ and $\mathcal{C} = \{(x_\alpha^c)\}_{c \in C}$ be nonempty sets of nets in a vector lattice E indexed by the same directed set A , and let $r \in \mathbb{R}$. The following holds.

i) If $\mathcal{B} \xrightarrow{c-o} 0$ and $\mathcal{C} \xrightarrow{c-o} 0$ then $r\mathcal{B} = \{(rx_\alpha^b)\}_{\alpha \in A; b \in B} \xrightarrow{c-o} 0$,

$$\text{sol}(\mathcal{B}) := \{(x_\alpha) : (\exists b \in B)(\forall \alpha \in A)|x_\alpha| \leq |x_\alpha^b|\} \xrightarrow{c-o} 0,$$

$\mathcal{B} \cup \mathcal{C} \xrightarrow{c-o} 0$, $\mathcal{B} + \mathcal{C} = \{(x_\alpha^b + x_\alpha^c) : b \in B, c \in C\} \xrightarrow{c-o} 0$, and $\text{co}(\mathcal{B}) \xrightarrow{c-o} 0$, where

$$\text{co}(\mathcal{B}) = \left\{ (x_\alpha) = \left(\sum_{k=1}^n r_k x_\alpha^{b_k} \right) : b_k \in B, r_k \in \mathbb{R}_+, \sum_{k=1}^n r_k = 1 \right\}.$$

ii) If $\mathcal{B} \xrightarrow{c-uo} 0$ and $\mathcal{C} \xrightarrow{c-uo} 0$ then $r\mathcal{B} \xrightarrow{c-uo} 0$, $\text{sol}(\mathcal{B}) \xrightarrow{c-uo} 0$, $\mathcal{B} \cup \mathcal{C} \xrightarrow{c-uo} 0$, $\mathcal{B} + \mathcal{C} \xrightarrow{c-uo} 0$, and $\text{co}(\mathcal{B}) \xrightarrow{c-uo} 0$.

iii) If $\mathcal{B} \xrightarrow{c-ru} 0$ and $\mathcal{C} \xrightarrow{c-ru} 0$ then $r\mathcal{B} \xrightarrow{c-ru} 0$, $\text{sol}(\mathcal{B}) \xrightarrow{c-ru} 0$, $\mathcal{B} \cup \mathcal{C} \xrightarrow{c-ru} 0$, $\mathcal{B} + \mathcal{C} \xrightarrow{c-ru} 0$, and $\text{co}(\mathcal{B}) \xrightarrow{c-ru} 0$.

Moreover, $x_\alpha \xrightarrow{o} 0 \iff \{(x_\alpha)\} \xrightarrow{c-o} 0$; $x_\alpha \xrightarrow{uo} 0 \iff \{(x_\alpha)\} \xrightarrow{c-uo} 0$; and $x_\alpha \xrightarrow{ru} 0 \iff \{(x_\alpha)\} \xrightarrow{c-ru} 0$.

Proof. i) Let $\mathcal{B} \xrightarrow{c-o} 0$ and $\mathcal{C} \xrightarrow{c-o} 0$. Take two nets $g_\beta \downarrow 0$ and $p_\gamma \downarrow 0$ in E such that, for every β and γ , there exist indices α_β and α_γ satisfying $|x_\alpha^b| \leq g_\beta$ for $\alpha \geq \alpha_\beta$ and $b \in B$, and $|x_\alpha^c| \leq p_\gamma$ for $\alpha \geq \alpha_\gamma$ and $c \in C$.

Obviously, $r\mathcal{B} \xrightarrow{c-o} 0$ and $\text{sol}(\mathcal{B}) \xrightarrow{c-o} 0$.

In order to show $\mathcal{B} \cup \mathcal{C} \xrightarrow{c-o} 0$, consider the net $(g_\beta + p_\gamma)_{\beta; \gamma} \downarrow 0$. Since $\mathcal{B} \cup \mathcal{C} = \{(x_\alpha^d)\}_{\alpha \in A; d \in B \cup C}$ then $|x_\alpha^d| \leq g_\beta + p_\gamma$ for $\alpha \geq \alpha_\beta, \alpha_\gamma$ and $d \in B \cup C$. For every β and γ find $\alpha_{(\beta, \gamma)} \geq \alpha_\beta, \alpha_\gamma$. Then $|x_\alpha^d| \leq g_\beta + p_\gamma$ for $\alpha \geq \alpha_{(\beta, \gamma)}$ and $d \in B \cup C$ as desired.

For proving $\mathcal{B} + \mathcal{C} \xrightarrow{c-o} 0$, consider $(g_\beta + p_\gamma)_{\beta; \gamma} \downarrow 0$, and pick $\alpha_{(\beta, \gamma)} \geq \alpha_\beta, \alpha_\gamma$. It follows $|x_\alpha^b + x_\alpha^c| \leq g_\beta + p_\gamma$ for $\alpha \geq \alpha_{(\beta, \gamma)}$, $b \in B$, and $c \in C$ as desired.

For proving $\text{co}(\mathcal{B}) \xrightarrow{c-o} 0$, let $b_1, \dots, b_n \in B$ and $r_1, \dots, r_n \in \mathbb{R}_+$ such that $\sum_{k=1}^n r_k = 1$. Since, for $\alpha \geq \alpha_\beta$ we have

$$\left| \sum_{k=1}^n r_k x_\alpha^{b_k} \right| \leq \sum_{k=1}^n r_k |x_\alpha^{b_k}| \leq \sum_{k=1}^n r_k g_\beta = g_\beta,$$

it follows $\text{co}(\mathcal{B}) \xrightarrow{c-o} 0$.

ii) It is a direct consequence of i).

iii) Let $\mathcal{B} \xrightarrow{c-ru} 0$ and $\mathcal{C} \xrightarrow{c-ru} 0$. Obviously, $r\mathcal{B} \xrightarrow{c-ru} 0$ and $\text{sol}(\mathcal{B}) \xrightarrow{c-ru} 0$.

Take $u, w \in E_+$ and two increasing sequences (α'_n) and (α''_n) of indices such that $|x_\alpha^b| \leq \frac{1}{n}u$ for $\alpha \geq \alpha'_n$ and $b \in B$, and $|x_\alpha^c| \leq \frac{1}{n}w$ for $\alpha \geq \alpha''_n$ and $c \in C$. For each n find some $\alpha_n \geq \alpha'_n, \alpha''_n$. Since $|x_\alpha^d| \leq \frac{1}{n}(u + w)$ for $\alpha \geq \alpha_n$ and $d \in B \cup C$ then $\mathcal{B} \cup \mathcal{C} \xrightarrow{c-ru} 0$.

Since $|x_\alpha^b + x_\alpha^c| \leq \frac{1}{n}(u + w)$ for all $\alpha \geq \alpha_n$, $b \in B$, and $c \in C$, it follows $\mathcal{B} + \mathcal{C} \xrightarrow{c\text{-ru}} 0$.

In order to show $\text{co}(\mathcal{B}) \xrightarrow{c\text{-ru}} 0$, let $b_1, \dots, b_n \in B$ and $r_1, \dots, r_n \in \mathbb{R}_+$ satisfy $\sum_{k=1}^n r_k = 1$. Since

$$\left| \sum_{k=1}^n r_k x_\alpha^{b_k} \right| \leq \sum_{k=1}^n r_k |x_\alpha^{b_k}| \leq \sum_{k=1}^n \frac{r_k}{n} u = \frac{1}{n} u$$

for all $\alpha \geq \alpha'_n$, we conclude $\text{co}(\mathcal{B}) \xrightarrow{c\text{-ru}} 0$.

The remaining part of the proof is obvious. □

Definition 2 gives rise to the following notions.

Definition 3. Let $\mathcal{T} \subseteq \mathcal{L}(E, F)$.

If E and F are vector lattices, the set \mathcal{T} is

- a) *collectively order continuous* (briefly, $\mathcal{T} \in \mathbf{L}_{oc}(E, F)$) if $\mathcal{T}(x_\alpha) = \{(Tx_\alpha)\}_{T \in \mathcal{T}} \xrightarrow{c\text{-o}} 0$ whenever $x_\alpha \xrightarrow{o} 0$.
- b) *collectively uo-continuous* (briefly, $\mathcal{T} \in \mathbf{L}_{uoc}(E, F)$) if $\mathcal{T}(x_\alpha) \xrightarrow{c\text{-uo}} 0$ whenever $x_\alpha \xrightarrow{uo} 0$.
- c) *collectively ru-continuous* (briefly, $\mathcal{T} \in \mathbf{L}_{rc}(E, F)$) if $\mathcal{T}(x_\alpha) \xrightarrow{c\text{-ru}} 0$ whenever $x_\alpha \xrightarrow{ru} 0$.
- d) *collectively order bounded* (briefly, $\mathcal{T} \in \mathbf{L}_{ob}(E, F)$) if the set $\mathcal{T}[0, b] = \{Tx : T \in \mathcal{T}; x \in [0, b]\}$ is order bounded for every $b \in E_+$.

If E is a vector lattice and F is a normed space, the set \mathcal{T} is

- e) *collectively order to norm bounded* (briefly, $\mathcal{T} \in \mathbf{L}_{omb}(E, F)$) if the set $\mathcal{T}[0, b]$ is norm bounded for every $b \in E_+$.

Clearly, an operator T is order bounded (order to norm bounded, o-, uo-, ru-continuous) iff the set $\{T\}$ is collectively order bounded (resp., collectively order to norm bounded, o-, uo-, ru-continuous). The following example shows that collectively o-bounded and/or collectively compact sets of o-continuous (or, uo-continuous) functionals need not to be o-continuous (uo-continuous) collectively.

Example 2.1. Consider a sequence (f_n) of functionals on ℓ^2

$$f_n \in (\ell^2)' : f_n(\mathbf{x}) = x_n,$$

for $\mathbf{x} = \sum_{k=1}^\infty x_k \mathbf{e}_k \in \ell^2$, where \mathbf{e}_k is the k -th unit vector of ℓ^2 . Then (f_n) is a collectively o-bounded collectively compact sequence of o- and uo-continuous functionals that is neither o- nor uo-continuous collectively.

2.2. On continuity of order to norm bounded operators. It

is well known that every order bounded operator from a BL to a NL is continuous (cf., [1, Theorem 1.31]). We extend this fact in the following strong collective way.

Theorem 1. *Every collectively order to norm bounded set of operators from a Banach lattice to a normed space is bounded in the operator norm.*

Proof. Let E be a BL, Y a NS, and $\mathcal{T} \in \mathbf{L}_{onb}(E, Y)$. Assume in contrary, \mathcal{T} is not bounded in the operator norm. Then the set $\mathcal{T}(B_E)$, and hence the set $\mathcal{T}((B_E)_+)$ is not norm bounded. Thus, there exist sequences (x_n) in $(B_E)_+$ and (T_n) in \mathcal{T} satisfying $\|T_n x_n\| \geq n^3$ for all $n \in \mathbb{N}$. Take $x := \|\cdot\| - \sum_{n=1}^{\infty} \frac{x_n}{n^2} \in E_+$. Since $\mathcal{T} \in \mathbf{L}_{onb}(E, Y)$, there exists $M \in \mathbb{R}_+$ such that $\mathcal{T}([0, x]) \subseteq MB_Y$. Since $\frac{x_n}{n^2} \in [0, x]$ then $\|T_n(\frac{x_n}{n^2})\| \leq M$ for every $n \in \mathbb{N}$. We get a contradiction:

$$M \geq \left\| T_n \left(\frac{x_n}{n^2} \right) \right\| \geq n \quad (\forall n \in \mathbb{N}).$$

Therefore, the set \mathcal{T} is norm bounded. □

It is worth noting the same argument combined with the Krein – Smulian theorem (cf., [5, Theorem 2.37]) proves norm boundedness of every collective order to norm bounded set of linear operators from an ordered Banach space with a closed generating cone to a normed space.

Corollary 1. *Every linear operator from a Banach lattice to a normed space that maps order intervals to norm bounded sets is bounded.*

2.3. Further elementary properties of collectively qualified sets.

Proposition 2. *Let E and F be vector lattices, and $r_1, r_2 \in \mathbb{R}$.*

- i) If \mathcal{T}_1 and \mathcal{T}_2 are nonempty subsets of $\mathbf{L}_{ob}(E, F)$, then $r_1\mathcal{T}_1 + r_2\mathcal{T}_2 = \{r_1T_1 + r_2T_2 : T_1 \in \mathcal{T}_1, T_2 \in \mathcal{T}_2\} \in \mathbf{L}_{ob}(E, F)$.*
- ii) If \mathcal{T}_1 and \mathcal{T}_2 are nonempty subsets of $\mathbf{L}_{oc}(E, F)$ (resp., $\mathbf{L}_{uoc}(E, F)$, $\mathbf{L}_{rc}(E, F)$) then $\mathcal{T}_1 \cup \mathcal{T}_2$ and $r_1\mathcal{T}_1 + r_2\mathcal{T}_2$ are subsets of $\mathbf{L}_{oc}(E, F)$ (resp., of $\mathbf{L}_{uoc}(E, F)$, $\mathbf{L}_{rc}(E, F)$).*

Proof. It follows directly from Proposition 1. □

For each element x of a VL E , we define the set $\mathcal{T}_x := \{y^\sim : |y| \leq |x|\}$ of functionals on the order dual E^\sim of E , where $y^\sim(f) = f(y)$ for each $f \in E^\sim = \mathcal{L}_{ob}(E, \mathbb{R})$.

Proposition 3. *Let E be a vector lattice. Then $\mathcal{T}_x \in \mathbf{L}_{oc}(E^\sim, \mathbb{R})$ for each $x \in E$.*

Proof. Let $x \in E$. In order to show that \mathcal{T}_x is collectively order continuous, let $f_\alpha \xrightarrow{o} 0$ in E^\sim and pick $g_\beta \downarrow 0$ in E^\sim so that, for each β there is an α_β with $|f_\alpha| \leq g_\beta$ for $\alpha \geq \alpha_\beta$. By the Riesz-Kantorovich formula, $|y^\sim(f_\alpha)| = |f_\alpha(y)| \leq |f_\alpha|(|y|) \leq |f_\alpha|(|x|) \leq g_\beta(|x|)$ for all $|y| \leq |x|$ and $\alpha \geq \alpha_\beta$. Since $g_\beta(|x|) \downarrow 0$, then $\mathcal{T}_x(f_\alpha) = \{(y^\sim(f_\alpha)) : |y| \leq |x|\} \xrightarrow{c-o} 0$. As $f_\alpha \xrightarrow{o} 0$ is arbitrary, we conclude $\mathcal{T}_x \in \mathbf{L}_{oc}(E^\sim, \mathbb{R})$. □

Let E^δ be a Dedekind completion of a VL E , F be a Dedekind complete VL, and $T \in (\mathcal{L}_{oc})_+(E, F)$. Define $T^\delta \in \mathcal{L}_{oc}(E^\delta, F)$ as the unique linear extension of an additive map $E_+ \ni y \rightarrow \sup_{y \geq x \in E} Tx$. The mapping

$(\mathcal{L}_{oc})_+(E, F) \ni T \rightarrow T^\delta$ has a unique extension to a Riesz isomorphism of $\mathcal{L}_{oc}(E, F)$ onto $\mathcal{L}_{oc}(E^\delta, F)$ (see, for example, [3, Theorem 1.84], [13, Theorem 3.2.3]).

Proposition 4. *Let a VL E be Archimedean, a VL F Dedekind complete, and $\mathcal{T} \subseteq \mathcal{L}_+(E, F)$. Then $\mathcal{T}^\delta \in \mathbf{L}_{oc}(E^\delta, F) \iff \mathcal{T} \in \mathbf{L}_{oc}(E, F)$.*

Proof. The regularity of E in E^δ (cf., [3, Theorems 1.23 and 1.41]) provides the implication $\mathcal{T}^\delta \in \mathbf{L}_{oc}(E^\delta, F) \implies \mathcal{T} \in \mathbf{L}_{oc}(E, F)$.

Now let $\mathcal{T} \in \mathbf{L}_{oc}(E, F)$ and $x_\alpha \xrightarrow{c-o} 0$ in E^δ . Then, there is $y_\beta \downarrow 0$ in E^δ such that, for each β there exists α_β with $|x_\alpha| \leq y_\beta$ for all $\alpha \geq \alpha_\beta$. Since E is majorizing in E^δ (cf., [3, Theorem 1.41]), we may assume that (y_β) is contained in E . Trivially, $y_\beta \downarrow 0$ in E . Since $\mathcal{T} \in \mathbf{L}_{oc}(E, F)$, we have $\{(Ty_\beta)\}_{T \in \mathcal{T}} \xrightarrow{c-o} 0$, and hence there is a net $g_\gamma \downarrow 0$ in F s.t., for each γ , there is β_γ with $|Ty_\beta| \leq g_\gamma$ for all $\beta \geq \beta_\gamma$ and $T \in \mathcal{T}$. As $\mathcal{T} \subseteq \mathcal{L}_+(E, F)$, we conclude $|T^\delta x_\alpha| \leq T^\delta |x_\alpha| \leq T^\delta y_{\beta_\gamma} = Ty_{\beta_\gamma} \leq g_\gamma$ for all $\alpha \geq \alpha_{\beta_\gamma}$ and $T \in \mathcal{T}$. Thus $\{(T^\delta x_\alpha)\}_{T \in \mathcal{T}} \xrightarrow{c-o} 0$, and hence $\mathcal{T}^\delta \in \mathbf{L}_{oc}(E^\delta, F)$. \square

2.4. Conditions for collectively order boundedness. It is long known that every o-continuous operator between VLs is order bounded (see, e.g., [3, Lemma 1.72]) whenever the order convergence is understood in the sense of [3, Definition 1.12] i.e., $x_\alpha \xrightarrow{o_1} x$ in a VL E if there exists a net $g_\alpha \downarrow 0$ in E s.t. $|x_\alpha - x| \leq g_\alpha$ for all α . Abramovich and Sirotkin proved in [2] that every o-continuous operator from an Archimedean VL to a VL is order bounded also if the order convergence is understood in the sense of Definition 1. The Archimedean assumption is essential in [2, Theorem 2.1] (see [11, Example 2.1]).

A short looking at the proof of [2, Theorem 2.1] tell us that Abramovich and Sirotkin proved indeed that every ru-continuous operator between arbitrary two VLs E and F is order bounded. Conversely, each $T \in \mathcal{L}_{ob}(E, F)$ is ru-continuous directly by Definition 1 c) because, for every $u \in E_+$ there is $w \in F$ with $T[-u, u] \subseteq [-w, w]$, and hence $|x_\alpha| \leq \frac{1}{n}u \implies |Tx_\alpha| \leq \frac{1}{n}w$.

The fact that $T \in \mathcal{L}_{ob}(E, F)$ iff T is ru-continuous was independently rediscovered by Taylor and Troitsky in [15, Theorem 5.1]. It is worth mentioned that this fact can also be derived from the nonstandard criteria of order boundedness [7, Theorem 1.7.2] (cf., [8, Theorem 4.6.2]).

We extend the Abramovich–Sirotkin–Taylor–Troitsky result as follows with the key idea of proof coming from [2].

Theorem 2. *Let E and F be vector lattices with E Archimedean. The following statements hold.*

i) Let $\mathcal{T} \subseteq \mathcal{L}(E, F)$. If $\mathcal{T}x_\alpha \xrightarrow{c-o} 0$ whenever $x_\alpha \downarrow 0$, then $\mathcal{T} \in \mathbf{L}_{ob}(E, F)$.

In particular, $\mathbf{L}_{oc}(E, F) \subseteq \mathbf{L}_{ob}(E, F)$.

ii) If $\dim(F) < \infty$ then $\mathbf{L}_{uoc}(E, F) \subseteq \mathbf{L}_{ob}(E, F)$.

Moreover, $\mathbf{L}_{ob}(E, F) = \mathbf{L}_{rc}(E, F)$ for all vector lattices E and F .

Proof. *i)* Let $[0, b]$ be an order interval in E . Like in the proof of [2, Theorem 2.1], let $\mathcal{I} = \mathbb{N} \times [0, b]$ be a set directed with the lexicographical order: $(m, z) \geq (n, y)$ iff either $m > n$ or $m = n$ and $z \geq y$. Let $x_{(k,y)} = \frac{1}{k}y \in [0, b]$. Since $0 \leq x_{(m,y)} = \frac{1}{m}y \leq \frac{1}{n}b = x_{(n,b)}$ for $(m, y) \geq (n, b)$ then $x_{(m,y)} \xrightarrow{ru} 0$, and hence $x_{(m,y)} \downarrow 0$ because E is Archimedean.

By the assumption, there exists a net $g_\beta \downarrow 0$ such that, for every β there exists (m_β, y_β) satisfying $|Tx_{(m,y)}| \leq g_\beta$ for all $(m, y) \geq (m_\beta, y_\beta)$ and $T \in \mathcal{T}$. Pick any g_{β_0} . Since $(m_{\beta_0} + 1, y) \geq (m_{\beta_0}, y_{\beta_0})$, it follows

$$\left| T\left(\frac{y}{m_{\beta_0} + 1}\right) \right| = |Tx_{(m_{\beta_0} + 1, y)}| \leq g_{\beta_0} \quad (y \in [0, b], T \in \mathcal{T}).$$

Then $|Ty| \leq (m_{\beta_0} + 1)g_{\beta_0}$ for $y \in [0, b]$ and $T \in \mathcal{T}$. Since $[0, b]$ is arbitrary, the set \mathcal{T} is collective order bounded.

ii) Let $[0, b] \subset E$. Take a directed set \mathcal{I} as in the proof of *i)*, and consider a net $x_{(k,y)} = \frac{1}{k}y$ in $[0, b]$. Then $x_{(m,y)} \xrightarrow{ru} 0$, and hence $x_{(m,y)} \xrightarrow{uo} 0$ as E is Archimedean. Because of $\dim(F) < \infty$, F has a strong unit, say w . Since $\mathcal{T} \in \mathbf{L}_{uoc}(E, F)$, there exists a net $g_\beta \downarrow 0$ in F such that, for every β , there exists (m_β, y_β) with $|Tx_{(m,y)}| \wedge w \leq g_\beta$ for all $(m, y) \geq (m_\beta, y_\beta)$ and $T \in \mathcal{T}$. As $\dim(F) < \infty$, by passing to a tail, we may assume $g_\beta \leq \frac{w}{2}$ for all β . So, $|Tx_{(m,y)}| \leq g_\beta$ for $(m, y) \geq (m_\beta, y_\beta)$ and $T \in \mathcal{T}$. Pick any β_0 . Since $(m_{\beta_0} + 1, y) \geq (m_{\beta_0}, y_{\beta_0})$,

$$\left| T\left(\frac{y}{m_{\beta_0} + 1}\right) \right| = |Tx_{(m_{\beta_0} + 1, y)}| \leq g_{\beta_0} \quad (y \in [0, b], T \in \mathcal{T}).$$

Thus, $|T(y)| \leq (m_{\beta_0} + 1)g_{\beta_0}$ for all $y \in [0, b]$ and $T \in \mathcal{T}$, and hence $\mathcal{T} \in \mathbf{L}_{ob}(E, F)$.

Now, let E and F be arbitrary vector lattices.

Let $\mathcal{T} \in \mathbf{L}_{ob}(E, F)$ and $x_\alpha \xrightarrow{ru} 0$. Then, for some $u \in E_+$, there exists an increasing sequence (α_n) of indices with $n x_\alpha \in [-u, u]$ for all $\alpha \geq \alpha_n$. Since $\mathcal{T} \in \mathbf{L}_{ob}(E, F)$ then $\mathcal{T}[-u, u] \subseteq [-w, w]$ for some $w \in F_+$, and hence $n\mathcal{T}x_\alpha \subseteq [-w, w]$ for $\alpha \geq \alpha_n$. It follows $\mathcal{T} \in \mathbf{L}_{rc}(E, F)$.

Finally, let $\mathcal{T} \in \mathbf{L}_{rc}(E, F)$ and $[0, b] \subset E$. Take a directed set \mathcal{I} and a net $x_{(k,y)} \xrightarrow{ru} 0$ as in the proof of *i)*. Then, for some $u \in E_+$ there exists an increasing sequence (k_n, y_n) in \mathcal{I} with $|T(\frac{y}{k})| = |Tx_{(k,y)}| \leq \frac{1}{n}u$ for all $(k, y) \geq (k_n, y_n)$ and $T \in \mathcal{T}$. In particular, $|T(\frac{y}{k_1+1})| = |Tx_{(k_1+1, y)}| \leq u$ for all $y \in [0, b]$ and all $T \in \mathcal{T}$. This implies that $|Ty| \leq (k_1 + 1)u$ for all $y \in [0, b]$ and all $T \in \mathcal{T}$. Since $[0, b]$ is arbitrary, $\mathcal{T} \in \mathbf{L}_{ob}(E, F)$. \square

It is an open question, whether the assumption $\dim(F) < \infty$ can be dropped in Theorem 2 *ii*). In favor of dropping this assumption, we have a relatively easy observation that each of non order bounded operators in the textbooks [3], [4], and [16] (see, [3, Exercise 30; p. 48], [4, Examples 2.38 and 4.73], [16, Exercise 98.7]) is not uo-continuous.

From the other hand, it is well known that a positive functional need not to be uo-continuous; e.g., $f \in (\ell^1)'$, $f(\mathbf{x}) = \sum_{k=1}^{\infty} x_k$, for $\mathbf{x} = \sum_{k=1}^{\infty} x_k \mathbf{e}_k \in \ell^1$, where \mathbf{e}_k is the k -th unit vector of ℓ^1 , is not uo-continuous as $f(\mathbf{e}_n) \equiv 1$ despite $\mathbf{e}_n \xrightarrow{uo} 0$ in ℓ^1 .

In the Banach lattice setting, Theorem 2 has the following application to extension of the well known fact that each o-continuous operator from a Banach lattice to a normed lattice is continuous.

Theorem 3. *Let $\mathcal{T} \in \mathbf{L}_{oc}(E, F)$, where E is a Banach lattice and F a normed lattice. Then the set \mathcal{T} is norm bounded.*

Proof. By Theorem 2, $\mathcal{T} \in \mathbf{L}_{ob}(E, F)$, and hence $\mathcal{T} \in \mathbf{L}_{onb}(E, F)$. An application of Theorem 1 completes the proof. \square

References

- [1] Y.A. Abramovich, C.D. Aliprantis, *An Invitation to Operator Theory*, Graduate Studies in Mathematics, Vol. 50, American Mathematical Society, Providence, RI, 2002. Zbl 1022.47001
- [2] Y. Abramovich, G. Sirotkin, *On order convergence of nets*, Positivity, **9**:3 (2005), 287–292. Zbl 1110.46002
- [3] C.D. Aliprantis, O. Burkinshaw, *Locally Solid Riesz Spaces with Applications to Economics*, 2nd ed., Math. Surveys and Monographs, Vol. 105, American Mathematical Society, Providence, RI, 2003. Zbl 1043.46003
- [4] C.D. Aliprantis and O. Burkinshaw, *Positive operators*, 2nd ed., Springer, Dordrecht, 2006. Zbl 1098.47001
- [5] C.D. Aliprantis, R. Tourky, *Cones and Duality*, Graduate Studies in Mathematics, Vol. 84, American Mathematical Society, Providence, RI, 2007. Zbl 1127.46002
- [6] P.M. Anselone, T.W. Palmer, *Collectively compact sets of linear operators*, Pac. J. Math., **25**:3 (1968), 417–422. Zbl 0157.45202
- [7] E.Yu. Emel'yanov, *Infinitesimal analysis and vector lattices*, Sib. Adv. Math., **6**:1 (1996), 19–70. Zbl 0854.46061
- [8] E.Y. Emelyanov, *Infinitesimals in vector lattices*, Chapter 4 in "Nonstandard Analysis and Vector Lattices", Mathematics and its Applications, Vol. 525, Kluwer Academic Publishers, Dordrecht – Boston – London, (2000), 161–230. Zbl 1001.46050
- [9] E. Emelyanov, *On collectively L -weakly compact sets of operators*, . Anal. Math., **51**:2 (2025), 447–455. Zbl 8079190.
- [10] E. Emelyanov, *On collectively σ -Levi sets of operators*, Vladikavkaz. Mat. Zh., **27**:1 (2025), 36–43. Zbl 1569.47049
- [11] E. Emelyanov, N. Erkurşun-Özcan, S. Gorokhova, *Collective Order Boundedness of Sets of Operators Between Ordered Vector Spaces*, Result. Math., **80**:3 (2025), paper no. 70. Zbl 8017038

- [12] S. Gorokhova, *On compact (limited) operators between Hilbert and Banach spaces*, *Filomat*, **38**:33 (2024), 11633–11637.
- [13] A. G. Kusraev, *Dominated Operators*, Mathematics and its Applications, Vol. 519, Kluwer Academic Publishers, Dordrecht – London, 2000. Zbl 0983.47025
- [14] P. Meyer-Nieberg, *Banach Lattices*, Springer–Verlag, Berlin – New York, 1991. Zbl 0743.46015
- [15] M.A. Taylor and V.G. Troitsky, *Bibasic sequences in Banach lattices*, *J. Funct. Anal.* **278**:10 (2020), Article ID 108448. Zbl 1445.46020
- [16] A.C. Zaanen, *Riesz Spaces. II*, North-Holland Mathematical Library, Vol. 30, North-Holland Publishing Co., Amsterdam – New York – Oxford, 1983. Zbl 0519.46001

EDUARD YURYEVIKH EMEL'YANOV
SOBOLEV INSTITUTE OF MATHEMATICS,
PR. KOPTYUGA, 4,
630090, NOVOSIBIRSK, RUSSIA
Email address: emelanov@math.nsc.ru